

Newton-Krylov generalized minimal residual algorithm in solving Volterra-Fredholm-Hammerstein integral equations

A. Zavvartorbati ¹

Abstract : In this paper, Galerkin and collocation methods have been applied on nonlinear Volterra-Fredholm-Hammerstein (VFH) integral equations, these methods are based on shifted Legendre polynomials, then methods transfer the finding solution of a nonlinear integral equation to finding the solution of nonlinear algebraic equations, in order to solve these nonlinear algebraic equations we use Newton method composed by generalized minimal residual (NGMRes) method, the iteration number and running time for implementation of NGMRes method are important parameters that have been considered to solve this type of integral equations. These methods are applied on several various nonlinear VFH integral equations that confirm accuracy and efficiency of the methods.

Keywords: Volterra-Fredholm-Hammerstein integral equations; Collocation method; Galerkin method; Spectral methods; Newton-Krylov GMRes; Shifted Legendre polynomials.

2020 Mathematics Subject Classification: 00A69; 64F12

Receive: 25 November 2020, **Accepted:** 15 January 2021

1 Introduction

Integral equations have found their applications in various fields of mathematics, sciences and technologies that have motivated a large amount of research works in recent years see [3, 13, 16, 20, 22, 32, 37–39]. VFH integral equations are one of the well-known nonlinear integral equations that arise in many fields including electricity and magnetism, communication theory, antenna synthesis problem, mathematical economics, population genetics, radiation, the particle transport problems of astrophysics and reactor theory, fluid mechanics, etc. The nonlinear VFH integral equations have been introduced in general form [28]

$$y(x) = f(x) + \lambda_1 \int_0^x k_1(x, t)G_1(t, y(t))dt + \lambda_2 \int_0^1 k_2(x, t)G_2(t, y(t))dt \quad 0 \leq x \leq 1, \quad (1.1)$$

where λ_1 and λ_2 are constants and $f(x)$ and kernels $k_1(x, t)$ and $k_2(x, t)$ are given functions assumed to have n th derivatives on intervals $0 \leq x, t \leq 1$, also $G_1(t, y(t))$ and $G_2(t, y(t))$ are given continuous functions which are nonlinear with respect to $y(t)$ and t . In [17] the existence of solution to nonlinear Hammerstein equations has been discussed.

Recently, many researchers have tried to solve this type of equations with several numerical methods and

¹Malek Ashtar University of Technology, Tehran, Iran. Email:zavvarahmad@gmail.com

algorithms, for example, a variation of the Nystrom method proposed in [19]. In [21] the nonlinear VFH integral equations have been solved via Hybrid of block-pulse functions. In [36] Haar wavelet collocation method for nonlinear VFH integral equations was proposed. In [10] Tau-collocation method has been applied on this type of integral equations and convergence of method has been investigated. In [40] Legendre Wavelet method have been applied. for more references see [1, 11, 12, 18].

Spectral methods are well-known methods that successfully applied in two past decades. These powerful tools are used to solve linear and nonlinear equations in many fields such as fluid dynamics, quantum dynamics and etc. Parand et al. in [26] discussed nonlinear Volterra-Fredholm Integro-Differential equations of Multi-Arbitrary Order. In [27] was solved Volterra's population growth model of arbitrary order using the generalized fractional order of the Chebyshev functions. In [29] was presented a new numerical algorithm based on the first kind of modified Bessel function to solve population growth in a closed system see [25, 31]. By applying spectral methods on nonlinear integral equations the problems are reduced to a system of nonlinear algebraic equations, so speed and accuracy of solving this nonlinear system have key role. Many works have been done to improve solving the nonlinear systems, Newton-Krylov algorithm is one of the most important and popular method that has been considered in many works such as [2, 7, 15].

We organized this paper as follows, in section 2, Legendre and shifted Legendre polynomials are explained, in section 3, Newton-Krylov GMRes algorithm will be discussed, in section 4, the methods are implemented on problem, section 5, will be devoted to convergence analysis, in section 6, the numerical results are reported, and at last section we will have a conclusion of our study.

2 Preliminary and notations

Properties of Legendre and shifted Legendre polynomials

Assuming that the Legendre polynomial of degree k is denoted by $P_k(z)$. The recurrence formulae of $P_k(z)$ is

$$P_{k+1}(z) = \frac{2k+1}{k+1}zP_k(z) - \frac{k}{k+1}P_{k-1}(z), \quad k = 1, 2, \dots,$$

$$P_0(z) = 1, \quad P_1(z) = z.$$

Substituting $z = 2x - 1$, Legendre polynomials are defined on the interval $[0, 1]$ that may be called shifted Legendre polynomials. The explicit analytical form of the shifted Legendre polynomial $P_k^*(x)$ of degree k is

$$P_k^*(x) = \sum_{j=0}^k (-1)^{j+k} \frac{(j+k)!x^j}{(k-j)!(k!)^2},$$

and this in turn, enables one to get

$$P_k^*(0) = (-1)^k, \quad P_k^*(1) = 1.$$

The orthogonality relation is

$$\int_0^1 P_i^*(x)P_j^*(x)dx = \begin{cases} \frac{1}{2j+1}, & \text{for } i = j, \\ 0, & \text{for } i \neq j. \end{cases}$$

Any square integrable function $y(x)$ defined on the interval $[0, 1]$, may be expressed in terms of shifted Legendre polynomials as

$$y(x) = \sum_{k=0}^{\infty} a_k F_k^*(x), \quad \text{where, } a_k = (2k+1) \int_0^1 y(x) F_k^*(x) dx, \quad k = 0, 1, \dots$$

3 Newton-Krylov algorithm

By applying spectral methods on nonlinear integral equations a system of nonlinear algebraic equations is obtained. Consider $F(x) = 0$ where $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a function $F(x) = (f_1(x), f_2(x), \dots, f_n(x))^T$ and $x \in \mathbb{R}^n$ is a vector. By applying the well-known Newton's iterative method:

$$F(x_{n+1}) = F(x_n) + (x_n - x_{n+1})F'(x_n),$$

where, $F'(x_n) = J(x_n)$ is the $n \times n$ Jacobian matrix. therefore

$$x_{n+1} = x_n - J(x_n)^{-1}F(x_n).$$

It is obvious in each iteration, a linear system must be solved, by increasing complexity of equations solving this linear system mostly could be time-consuming process. one of the best ideas to overcome this problem is to use NGMRes.

A framework for GMRes implementation

Begin

1. $r = b - Ax$, $v_1 = r/\|r\|_2$, $\rho = \|r\|_2$, $\beta = \rho$, $k = 0$.
 2. **while** $\rho > \epsilon\|b\|_2$ and $k < k_{max}$ **do**
 - 2.a $k = k + 1$.
 - 2.b Apply Arnoldi to obtain H_k and V_{k+1} from V_k and H_{k-1} .
 - 2.c $e_1 = (1, 0, \dots, 0)^T \in \mathbb{R}^{k+1}$.
 - 2.d Solve $\min \|\beta e_1 - H_k y_k\|_{\mathbb{R}^{k+1}}$ for $y_k \in \mathbb{R}^k$.
 - 2.e $\rho = \|\beta e_1 - H_k y_k\|_{\mathbb{R}^{k+1}}$.
 - 2.f $x_k = x_0 + V_k y_k$
- end while**

End

For more details see[8, 9, 33].

4 Proposed methods

Spectral methods belong to the family of weighted residual methods (WRMs). WRMs approximate the solution $y(x)$ of the integral equation (1.1) by a finite sum

$$y(x) \simeq y_N(x) = \sum_{k=0}^N a_k \phi_k(x),$$

that $\phi_k(x)$, is trial function, and a_k , $k = 0..N$ are unknown coefficients, by substitution $y_N(x)$ in Eq. (1.1), leads us to residual function

$$Res_N(x) = f(x) + \lambda_1 \int_0^x k_1(x,t)G_1(t, y_N(t))dt + \lambda_2 \int_0^1 k_2(x,t)G_2(t, y_N(t)) - y_N(x)dt. \quad (4.1)$$

The notion of the WRMs is to force the residual to zero in suitable norm,

$$\langle Res_N(x), \Psi_j \rangle_{\omega} = \int_{\Omega} Res_N(x) \Psi_j(x) \omega(x) dx = 0, \quad 0 \leq j \leq N \quad (4.2)$$

which Ψ_j is the test function, and ω is a weight function. Definition of shifted Legendre collocation method and shifted Legendre Galerkin method is as follows in table 1:

where $P_i^*(x)$, is shifted Legendre of degree i , and $\delta(x - x_j)$ is Dirac delta function and also x_j is j th zero

Table 1: Definition of shifted Legendre collocation method and shifted Legendre Galerkin method.

Method	Weight function	Trial function	Test function
Shifted Legendre collocation method	1	$P_i^*(x)$	$\delta(x - x_j)$
Shifted Legendre Galerkin method	1	$P_i^*(x)$	$P_j^*(x)$

of shifted Legendre of degree $N+1$. To calculate the Eq. (4.2) we use the Gauss-Legendre integration rule [5]. Finally we generate $N + 1$ nonlinear equations with $N + 1$ unknown coefficients, to solve this system of nonlinear algebraic equations and obtaining coefficients $a_k, k = 0, \dots, N$ we have used NGMRes method.

5 Error estimate

In this section, the theorem on convergence analysis and error estimation of the proposed is discussed. Suppose $y(x)$ is sufficiently smooth function on $[0,1]$, and $p_n(x)$ is the interpolating polynomial to y at x_i points, where $x_i, i = 0, \dots, N$ are the roots of chebyshev polynomial of degree $N + 1$ in $[0, 1]$, and then

$$y(x) - p_n(x) = \frac{y^{(N+1)}(\xi)}{(N+1)!} \prod_{i=0}^N (x - x_i) \quad \xi \in [0, 1], \quad (5.1)$$

then

$$|y(x) - p_n(x)| \leq \frac{M_N(1)^{N+1}}{2^{2N+1}(N+1)!}, \quad (5.2)$$

where, $M_N = \max_{0 \leq x \leq 1} |y^{(N+1)}(x)|$.

Theorem 5.1. Suppose that $y_N(x) = A^T P(x)$ be the shifted Legendre polynomials expansion of the exact solution, $y(x)$, where

$$A = [a_0, a_1, \dots, a_N]^T,$$

and

$$P(x) = [P_0(x), P_1(x), \dots, P_N(x)]^T.$$

Let $\bar{y}_N(x) = \sum_{i=0}^N \bar{a}_i P_i(x)$ be the approximate solution obtained by the method proposed in Section 4 and $M_N = \max_{0 \leq x \leq 1} |y^{N+1}(x)|$, then, there exist real numbers α and β_N such that

$$\|y(x) - \bar{y}_N(x)\|_2 \leq \alpha \frac{M_N(1)^{N+1}}{2^{2N+1}(N+1)!} + \beta_N \|A - \bar{A}\|_2, \quad (5.3)$$

where

$$\bar{A} = [\bar{a}_0, \bar{a}_1, \dots, \bar{a}_N]^T,$$

also the norm on the right hand side is the Euclidian norm for vectors.

Proof. Let $R_n[x]$ be the space of all real-valued polynomials of degree $\leq n$. Using the definition, $y_n(x)$ and $\bar{y}_n(x)$ are in $R_n[x]$ and $y_n(x)$ is the best approximation of $y(x)$ in $R_n[x]$. It's clear that

$$\|y(x) - \bar{y}_N(x)\|_2 \leq \|y(x) - y_N(x)\|_2 + \|y_N(x) - \bar{y}_N(x)\|_2. \quad (5.4)$$

By using Eq. (5.2) we have

$$\begin{aligned} \|y(x) - y_N(x)\|_2 &= \left(\int_0^l |y(x) - y_N(x)|^2 dx \right)^{\frac{1}{2}} \\ &\leq \left(\int_0^l \left[\frac{M_N(1)^{N+1}}{2^{2N+1}(N+1)!} \right]^2 dx \right)^{\frac{1}{2}} \\ &= \sqrt{l} \frac{M_N(1)^{N+1}}{2^{2N+1}(N+1)!}. \end{aligned} \quad (5.5)$$

Also, we have

$$\begin{aligned} \|y_N(x) - \bar{y}_N(x)\|_2 &= \left(\int_0^l \left[\sum_{i=0}^N (a_i - \bar{a}_i) P_i(x) \right]^2 dx \right)^{\frac{1}{2}} \\ &\leq \left(\int_0^l \left[\sum_{i=0}^N |a_i - \bar{a}_i|^2 \right] \left[\sum_{i=0}^N |P_i(x)|^2 \right] dx \right)^{\frac{1}{2}} \\ &= \left[\sum_{i=0}^N |a_i - \bar{a}_i|^2 \right]^{\frac{1}{2}} \left(\sum_{i=0}^N \int_0^l |P_i(x)|^2 dx \right)^{\frac{1}{2}} \\ &= \|A - \bar{A}\|_2 \left(l \sum_{i=0}^N \frac{1}{2i+1} \right)^{\frac{1}{2}}. \end{aligned} \quad (5.6)$$

Finally from Eqs. (5.4)-(5.6) we conclude that Eq. (5.3) is valid with

$$\alpha = \sqrt{l}, \quad \beta_N = \sqrt{l \left(\sum_{i=0}^N \frac{1}{2i+1} \right)}.$$

□

6 Illustrative Examples

In this section, In order to illustrate the performance of presented methods in solving VFH integral equations and justify the efficiency of the methods, we considered some various examples. To study the convergence behaviour of the methods, the maximum and mean errors have been used with the following definition, respectively

$$\begin{aligned} \|e_{(N)}\|_{\infty} &= \max_{(x) \in [0,1]} |u_{ex}(x) - u_N(x)| \\ \|e_{(N)}\|_2 &= \left(\int_0^1 (u_{ex}(x) - u_N(x))^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

The numerical implementation is carried out in Microsoft.maple.15, with hardware configuration: desktop 32-bit Intel core 2 Due CPU, 4 GB of RAM, 32-bit operation system.

Example 6.1. Consider nonlinear VFH integral equation given in [28], with the exact solution of $y(x)=\cos(x)$:

$$y(x) = 2 \cos(x) - 2 + 3 \int_0^x \sin(x-t)y^2(t) dt + \frac{6}{7-6 \cos(1)} \int_0^1 (1-t) \cos^2(x)(t+y(t)) dt, \quad (6.1)$$

we solve the Eq. (6.1), Table 2 shows the $\|e\|_2$, $\|e\|_\infty$, running time and number of iteration for implementation of Newton-Krylov algorithm, in Table 3 and Table 4, the absolute error for collocation and Galerkin methods at some point for different number of N is shown, respectively, Figure 1 displays the absolute error for $N = 15$.

Table 2: Some numerical results for example 6.1

N	Collocation method				Galerkin Method			
	$\ e\ _2$	$\ e\ _\infty$	Time (s)	Iteration	$\ e\ _2$	$\ e\ _\infty$	Time (s)	Iteration
2	3.71×10^{-3}	1.05×10^{-2}	1.21	3	4.82×10^{-3}	9.19×10^{-3}	3.26	2
4	2.53×10^{-6}	4.82×10^{-6}	1.74	5	2.12×10^{-6}	4.81×10^{-6}	5.43	3
6	3.67×10^{-9}	6.12×10^{-9}	1.86	5	3.92×10^{-9}	6.56×10^{-9}	6.07	3
8	2.29×10^{-12}	4.49×10^{-12}	2.15	5	2.74×10^{-12}	4.95×10^{-12}	7.18	5
10	1.57×10^{-15}	2.75×10^{-15}	3.01	5	6.89×10^{-14}	1.52×10^{-13}	7.93	5

Table 3: Absolute error of collocation method for example 6.1

X	N=2	N=4	N=6	N=8	N=10
0.0	1.02×10^{-4}	7.18×10^{-7}	5.06×10^{-10}	7.41×10^{-14}	3.17×10^{-17}
0.2	2.35×10^{-4}	7.80×10^{-7}	8.99×10^{-11}	8.22×10^{-14}	5.65×10^{-17}
0.4	8.86×10^{-4}	8.21×10^{-7}	6.45×10^{-10}	9.35×10^{-13}	1.27×10^{-16}
0.6	1.17×10^{-3}	9.00×10^{-7}	6.90×10^{-10}	1.00×10^{-12}	2.10×10^{-16}
0.8	1.04×10^{-3}	2.15×10^{-6}	9.32×10^{-10}	1.09×10^{-12}	2.93×10^{-16}
1.0	1.13×10^{-3}	2.80×10^{-6}	1.20×10^{-9}	1.31×10^{-12}	3.68×10^{-16}
$\ e\ _2$	3.71×10^{-3}	2.53×10^{-6}	3.67×10^{-9}	2.29×10^{-12}	1.57×10^{-15}

Example 6.2. Consider nonlinear VFH integral equation given in [28], with the exact solution of

$$y(x) = \ln\left(\frac{\theta^2}{2 \cos^2\left(\frac{\theta}{2}\left(x - \frac{1}{2}\right)\right)}\right), \theta = 4 \cos\left(\frac{\theta}{4}\right) :$$

$$y(x) = \int_0^1 K(x,t)e^{y(t)} dt, \quad (6.2)$$

where

$$K(x,t) = \begin{cases} -t(1-x), & 0 \leq t \leq x; \\ -x(1-t), & x \leq t \leq 1. \end{cases} \quad (6.3)$$

Table 4: Absolute error of Galerkin method for example 6.1

X	N=2	N=4	N=6	N=8	N=10
0.0	2.22×10^{-4}	8.40×10^{-8}	6.61×10^{-10}	4.37×10^{-13}	2.93×10^{-14}
0.2	5.81×10^{-4}	2.66×10^{-7}	8.47×10^{-10}	8.75×10^{-13}	4.98×10^{-14}
0.4	7.36×10^{-4}	8.87×10^{-7}	2.28×10^{-10}	1.04×10^{-12}	1.11×10^{-15}
0.6	9.25×10^{-4}	1.16×10^{-6}	9.75×10^{-10}	1.91×10^{-12}	2.30×10^{-15}
0.8	2.07×10^{-3}	9.98×10^{-7}	1.03×10^{-9}	1.40×10^{-12}	2.06×10^{-14}
1.0	3.36×10^{-3}	2.43×10^{-6}	2.05×10^{-9}	2.58×10^{-12}	1.32×10^{-14}
$\ e\ _2$	4.82×10^{-3}	2.12×10^{-6}	3.92×10^{-9}	2.74×10^{-12}	6.89×10^{-14}

we solve the Eq. (6.2), Table 5 shows the $\|e\|_2$, $\|e\|_\infty$, running time and number of iteration for implementation of Newton-Krylov algorithm, in Table 6 and Table 7, the absolute error for collocation and Galerkin methods at some point for different number of N is shown, respectively. Figure 2 displays the absolute error of collocation and Galerkin methods for $N = 25$ and $N = 20$, respectively.

Table 5: Some numerical results for example 6.2

N	Collocation method				Galerkin Method			
	$\ e\ _2$	$\ e\ _\infty$	Time (s)	Iteration	$\ e\ _2$	$\ e\ _\infty$	Time (s)	Iteration
5	3.12×10^{-5}	5.45×10^{-5}	1.16	16	5.27×10^{-9}	9.29×10^{-9}	3.58	5
10	4.96×10^{-8}	1.20×10^{-8}	1.53	17	4.80×10^{-14}	7.17×10^{-14}	26.10	3
15	1.22×10^{-12}	1.81×10^{-11}	1.28	23	9.39×10^{-15}	1.86×10^{-14}	32.78	3
20	3.17×10^{-14}	6.69×10^{-14}	3.40	30	8.11×10^{-26}	1.42×10^{-25}	52.69	5
25	6.18×10^{-18}	1.75×10^{-18}	4.41	35	-	-	-	-

Table 6: Absolute error of collocation method for example 6.2

X	N=5	N=10	N=15	N=20	N=25
0.0	1.91×10^{-6}	7.52×10^{-9}	8.78×10^{-13}	4.82×10^{-15}	1.52×10^{-19}
0.2	2.85×10^{-6}	4.99×10^{-9}	9.83×10^{-13}	9.07×10^{-15}	5.33×10^{-19}
0.4	6.62×10^{-6}	7.55×10^{-9}	9.14×10^{-13}	2.94×10^{-15}	1.27×10^{-16}
0.6	9.66×10^{-6}	3.49×10^{-9}	8.10×10^{-13}	5.29×10^{-15}	3.84×10^{-19}
0.8	4.53×10^{-6}	9.14×10^{-9}	1.13×10^{-12}	2.38×10^{-15}	2.57×10^{-19}
1.0	7.98×10^{-6}	1.02×10^{-8}	1.27×10^{-12}	8.75×10^{-15}	4.25×10^{-19}
$\ e\ _2$	3.12×10^{-5}	1.27×10^{-8}	1.22×10^{-12}	3.17×10^{-14}	6.18×10^{-18}

Example 6.3. Consider nonlinear Volterra-Hammerstein integral equation given in [36], with initial conditions $y(0)=1$. which has the exact solution $y(x)=exp(-x)$:

$$y(x) = \frac{3}{2} - \frac{1}{2}exp(-2x) - \int_0^x (y(t)^2 + y(t)) dt, \tag{6.4}$$





Figure 1: Absolute error of methods for Example 6.1

we solve the Eq. (6.4), Table 8 shows the $\|e\|_2$, $\|e\|_\infty$, running time and number of iteration for implementation of Newton-Krylov algorithm, in Table 9 and Table 10, the absolute error for collocation and Galerkin methods at some point for different number of N is shown, respectively, Figure 3 displays the

Table 7: Absolute error of Galerkin method for example6.2

X	N=5	N=10	N=15	N=20
0.0	3.90×10^{-10}	6.27×10^{-15}	6.45×10^{-17}	4.13×10^{-27}
0.2	3.22×10^{-10}	9.07×10^{-15}	5.39×10^{-16}	8.67×10^{-27}
0.4	9.39×10^{-10}	9.68×10^{-15}	9.74×10^{-17}	7.77×10^{-27}
0.6	4.32×10^{-10}	9.55×10^{-15}	9.45×10^{-17}	4.01×10^{-27}
0.8	2.49×10^{-10}	1.53×10^{-15}	3.97×10^{-16}	1.38×10^{-26}
1.0	1.28×10^{-10}	2.70×10^{-15}	4.49×10^{-16}	6.40×10^{-27}
$\ e\ _2$	5.27×10^{-9}	4.80×10^{-14}	9.39×10^{-15}	8.11×10^{-26}

absolute error for $N = 15$.

Table 8: Some numerical results for example 6.3

N	Collocation method				Galerkin Method			
	$\ e\ _2$	$\ e\ _\infty$	Time (s)	Iteration	$\ e\ _2$	$\ e\ _\infty$	Time (s)	Iteration
2	6.49×10^{-4}	1.67×10^{-3}	0.52	2	4.17×10^{-4}	8.59×10^{-4}	1.26	3
4	2.72×10^{-6}	4.84×10^{-6}	1.04	3	1.10×10^{-6}	2.22×10^{-6}	1.60	5
6	2.03×10^{-9}	4.61×10^{-9}	1.29	3	1.30×10^{-9}	2.55×10^{-9}	1.56	3
8	2.18×10^{-11}	4.22×10^{-11}	1.37	3	8.03×10^{-13}	1.87×10^{-12}	1.92	5
10	8.96×10^{-16}	1.81×10^{-15}	1.42	5	3.18×10^{-16}	9.46×10^{-16}	2.17	5


Table 9: Absolute error of collocation method for example 6.3

X	N=2	N=4	N=6	N=8	N=10
0.0	4.04×10^{-5}	3.15×10^{-7}	5.49×10^{-10}	9.20×10^{-13}	7.08×10^{-17}
0.2	8.11×10^{-5}	2.89×10^{-6}	8.68×10^{-10}	8.39×10^{-12}	7.88×10^{-17}
0.4	8.86×10^{-5}	9.95×10^{-7}	1.12×10^{-9}	4.25×10^{-12}	9.16×10^{-17}
0.6	6.81×10^{-5}	1.84×10^{-6}	8.49×10^{-10}	8.14×10^{-12}	8.11×10^{-17}
0.8	4.39×10^{-5}	9.32×10^{-7}	7.51×10^{-10}	9.15×10^{-12}	9.97×10^{-17}
1.0	9.66×10^{-5}	7.20×10^{-7}	6.31×10^{-10}	5.48×10^{-12}	9.56×10^{-17}
$\ e\ _2$	6.49×10^{-4}	2.72×10^{-6}	2.03×10^{-9}	2.18×10^{-11}	8.96×10^{-16}


Example 6.4. Consider nonlinear Volterra-Hammerstein integral equation given in [12], with the exact solution of $y(x)=x$:

$$y(x) = x + \frac{1}{x+1}(e^{-(x+1)} - 1) + \int_0^x \frac{1}{x} e^{-\frac{t}{x}} e^{-y(t)} dt. \tag{6.5}$$

we solve the Eq. (6.5), Table 11 shows the $\|e\|_2$, $\|e\|_\infty$, running time and number of iteration for implementation of Newton-Krylov algorithm, in Table 12 and Table 13, the absolute error for collocation



(a) Collocation method



(b) Galerkin method

Figure 2: Absolute error of methods for Example 5.2.

and Galerkin methods at some point for different number of N is shown, respectively, Figure 4 displays the absolute error for $N = 15$.

Table 10: Absolute error of Galerkin method for example 6.3

X	N=2	N=4	N=6	N=8	N=10
0.0	8.71×10^{-5}	1.35×10^{-7}	9.75×10^{-11}	6.08×10^{-15}	5.41×10^{-17}
0.2	7.66×10^{-5}	9.27×10^{-8}	1.33×10^{-10}	1.05×10^{-14}	1.78×10^{-17}
0.4	8.51×10^{-5}	8.61×10^{-8}	6.05×10^{-10}	1.28×10^{-14}	2.42×10^{-17}
0.6	6.45×10^{-5}	2.47×10^{-7}	9.75×10^{-10}	2.47×10^{-14}	1.38×10^{-17}
0.8	8.17×10^{-5}	7.75×10^{-7}	8.30×10^{-10}	5.15×10^{-14}	1.25×10^{-17}
1.0	6.76×10^{-5}	6.15×10^{-7}	5.72×10^{-10}	4.96×10^{-14}	2.07×10^{-17}
$\ e\ _2$	4.17×10^{-4}	1.10×10^{-6}	1.30×10^{-9}	8.03×10^{-13}	3.18×10^{-16}

Table 11: Some numerical results for example 6.4


N	Collocation method				Galerkin Method			
	$\ e\ _2$	$\ e\ _\infty$	Time (s)	Iteration	$\ e\ _2$	$\ e\ _\infty$	Time (s)	Iteration
2	8.03×10^{-20}	1.59×10^{-19}	0.29	6	6.50×10^{-18}	1.16×10^{-17}	0.21	6
4	1.09×10^{-24}	2.33×10^{-24}	0.31	9	2.03×10^{-28}	2.42×10^{-28}	0.27	7
6	8.73×10^{-42}	1.63×10^{-41}	0.47	17	2.03×10^{-43}	2.19×10^{-43}	0.36	7
8	1.12×10^{-58}	2.28×10^{-58}	0.52	23	7.65×10^{-48}	1.05×10^{-47}	0.42	7
10	1.27×10^{-63}	2.42×10^{-63}	1.07	27	3.07×10^{-56}	6.01×10^{-56}	0.44	8

Table 12: Absolute error of collocation method for example 6.4


X	N=2	N=4	N=6	N=8	N=10
0.0	6.08×10^{-22}	6.12×10^{-25}	4.98×10^{-43}	3.05×10^{-59}	8.59×10^{-65}
0.2	7.71×10^{-22}	3.52×10^{-25}	5.09×10^{-43}	2.31×10^{-59}	9.10×10^{-65}
0.4	8.23×10^{-21}	6.19×10^{-25}	5.36×10^{-43}	3.81×10^{-59}	2.46×10^{-64}
0.6	3.35×10^{-21}	8.72×10^{-25}	3.61×10^{-43}	4.76×10^{-59}	4.85×10^{-64}
0.8	1.91×10^{-20}	9.12×10^{-25}	6.41×10^{-43}	2.69×10^{-59}	3.35×10^{-64}
1.0	1.87×10^{-20}	9.90×10^{-25}	4.23×10^{-43}	7.88×10^{-59}	5.07×10^{-64}
$\ e\ _2$	8.03×10^{-20}	1.09×10^{-24}	8.73×10^{-42}	1.12×10^{-58}	1.27×10^{-63}

Table 13: Absolute error of Galerkin method for example 6.4

X	N=2	N=4	N=6	N=8	N=10
0.0	4.20×10^{-19}	9.14×10^{-30}	8.52×10^{-45}	8.02×10^{-50}	9.93×10^{-58}
0.2	7.02×10^{-19}	9.96×10^{-30}	8.68×10^{-45}	7.20×10^{-50}	2.49×10^{-57}
0.4	4.17×10^{-19}	2.40×10^{-29}	9.92×10^{-45}	1.23×10^{-49}	2.53×10^{-57}
0.6	3.52×10^{-19}	2.37×10^{-29}	1.38×10^{-44}	1.17×10^{-49}	3.48×10^{-57}
0.8	9.44×10^{-19}	4.29×10^{-29}	2.42×10^{-44}	1.28×10^{-49}	2.28×10^{-57}
1.0	7.84×10^{-19}	5.50×10^{-29}	5.49×10^{-44}	3.43×10^{-49}	4.12×10^{-57}
$\ e\ _2$	6.50×10^{-18}	2.03×10^{-28}	2.03×10^{-43}	7.65×10^{-48}	3.07×10^{-56}




(a) Collocation method




(b) Galerkin method

Figure 3: Absolute error of methods for Example 6.3



(a) Collocation method



(b) Galerkin method

Figure 4: Absolute error of methods for Example 6.4

7 Conclusion

In this paper, shifted Legendre collocation method and shifted Legendre Galerkin method were used to solve nonlinear VFH integral equations. By applying these two methods on nonlinear VFH integral equations the problems are reduced to a system of nonlinear algebraic equations, therefore, running time and accuracy in solving this system is an important part to approximate nonlinear VFH integral equations. In this work NGMRes method was used to solve obtained nonlinear algebraic equations system . The comparison of the obtained results of proposed methods and exact solutions shows that present methods are powerful tools to find the numerical solutions of such equations.

References

- [1] M.A. Abdou, Khamis I. Mohamed, A.S. Ismail, On the numerical solutions of FredholmVolterra integral equation, *Applied mathematics and computation*, 146(2-3) 2003, 713-728.
- [2] H. Asgharzadeh, I. Borazjani, A NewtonKrylov method with an approximate analytical Jacobian for implicit solution of NavierStokes equations on staggered overset-curvilinear grids with immersed boundaries, *Journal of computational physics*, 331 2017, 227-256.
- [3] J. Boersma and E. Danick, On the solution of an integral equation arising in potential problems for circular and elliptic disks, *SIAM Journal on Applied Mathematics*, 53(4) 1993, 931-941.
- [4] J.P. Boyd, *Chebyshev and Fourier spectral methods*, 2nd ed, New York Dover, 2000.
- [5] R.L. Burden, J.D. Faires, *Numerical Analysis*, Youngstown State University, Youngstown, 2001.
- [6] Y. Chen, C. Shen, A Jacobian-free Newton-GMRES (m) method with adaptive preconditioner and its application for power flow calculations, *IEEE Transactions on Power Systems*, 21(3) 2006, 1096-1103.
- [7] J. Chen, C. Vuik, Globalization technique for projected NewtonKrylov methods, *International Journal for Numerical Methods in Engineering*, 110(7) 2017, 661-674.
- [8] D. Fadrani, V. Rostami, K. Maleknejad, Fast iterative methods for solving of boundary nonlinear integral equations with singularity, *Journal of Computational Analysis and Applications*, 1(2) 1999, 219-234.
- [9] D. Fadrani, V. Rostami, K. Maleknejad, Preconditioners for solving stochastic boundary integral equations with weakly singular kernels, *Computing*, 63(1) 1999, 47-67.
- [10] Z. Gouyandeh, T. Allahviranloo, A. Armand, Numerical solution of nonlinear VolterraFredholmHammerstein integral equations via Tau-collocation method with convergence analysis, *Journal of Computational and Applied Mathematics* 100(308) 2016, 435-446.
- [11] M. Hadizadeh, R. Azizi, A reliable computational approach for approximate solution of Hammerstein integral equations of mixed type, *International Journal of Computer Mathematics* 81(7) 2004, 889-900.
- [12] M. Hadizadeh, M. Mohamadsohi, Numerical solvability of a class of Volterra-Hammerstein integral equations with noncompact kernels, *Journal of Applied Mathematics*, 2005(2) 2005, 171-181.
- [13] S. Hatamzadeh-Varmazyar, M. Naser-Moghadasi, E. Babolian, Z. Masouri, Numerical approach to survey the problem of electromagnetic scattering from resistive strips based on using a set of orthogonal basis functions, *Progress In Electromagnetics Research*, 81 2008, 393-412.

- [14] G. Han, Asymptotic error expansion variation of a collocation method for Volterra-Hammerstein equations, *Appl. Numer. Math.*, 13 1993, 357-369.
- [15] D.A. Knoll, D.E. Keyes, Jacobian-free Newton-Krylov methods: a survey of approaches and applications, *Journal of Computational Physics*, 193(2) 2004, 357-397.
- [16] E.V. Kovalenko, Some approximate methods of solving integral equations of mixed problems, *Journal of Applied Mathematics and Mechanics*, 53(1) 1989, 85-92.
- [17] F. Li, Y. Li, Z. Liang, Existence of solutions to nonlinear Hammerstein integral equations and applications, *Journal of Mathematical Analysis and Applications*, 323(1) 2006, 209-227.
- [18] M. Lakestani, M. Razzaghi, M. Dehghan, Solution of nonlinear Fredholm-Hammerstein integral equations by using semiorthogonal spline wavelets, *Mathematical problems in engineering*, 2005(1) 2005, 113-121.
- [19] L.J. Lardy, A Variation of Nyström's Method for Hammerstein Equations, *The Journal of Integral Equations*, 1981, 43-60.
- [20] A.V. Manzhirov, A mixed integral equation of mechanics and a generalized projection method of its solution, *Doklady Physics*, 61(10) 2016.
- [21] H.R. Marzban, H.R. Tabrizidooz, M. Razzaghi, A composite collocation method for the nonlinear mixed Volterra-Fredholm-Hammerstein integral equations, *Communications in Nonlinear Science and Numerical Simulation*, 16(3) 2011, 1186-1194.
- [22] M.V. Mirkin and A.J. Bard, Multidimensional integral equations: a new approach to solving micro-electrode diffusion problems: Part 2. Applications to microband electrodes and the scanning electrochemical microscope, *Journal of Electroanalytical Chemistry*, 323(1-2) 1992, 29-51.
- [23] Y. Ordokhani, Solution of nonlinear Volterra-Fredholm-Hammerstein integral equations via rationalized Haar functions, *Applied Mathematics and Computation*, 180(2) 2006, 436-443.
- [24] Y. Ordokhani, Solution of Fredholm-Hammerstein integral equations with Walsh-Hybrid functions, *International Mathematical Forum*, 4(20) 2009.
- [25] K. Parand, A. Bahramnezhad, H. Farahani, A numerical method based on rational Gegenbauer functions for solving boundary layer flow of a Powell-Eyring non-Newtonian fluid, *Computational and Applied Mathematics*, 37(5) 2018, 6053-6075
- [26] K. Parand, M. Delkhosh, Operational matrices to solve nonlinear volterra-fredholm integro-differential equations of multi-arbitrary order, *Gazi University Journal of Science*, 29(4) 2016, 895-907.
- [27] K. Parand, M. Delkhosh, Solving Voltterra's population growth model of arbitrary order using the generalized fractional order of the Chebyshev functions Authors, *Ricerche di Matematica*, 65(1) 2016, 307-328.
- [28] K. Parand, J.A. Rad, Numerical solution of nonlinear Volterra-Fredholm-Hammerstein integral equations via collocation method based on radial basis functions, *Applied Mathematics and Computation*, 218(9) 2012, 5292-5309.
- [29] K. Parand, J.A. Rad, M. Nikarya, A new numerical algorithm based on the first kind of modified Bessel function to solve population growth in a closed system, *International Journal of Computer Mathematics*, 91(6) 2014, 1239-1254.

- [30] K. Paranda, M. Nikarya, A numerical method to solve the 1D and the 2D reaction diffusion equation based on Bessel functions and Jacobian free Newton-Krylov subspace methods, *The European Physical Journal Plus* 132(11) 2017, 1-18.
- [31] K. Parand, S. Latifi, M.M. Moayeri, M. Delkhosh, Generalized Lagrange Jacobi Gauss-Lobatto (GLJGL) Collocation Method For Solving Linear and Nonlinear Fokker-Planck equations, *Communications in Theoretical Physics*, 69(5) 2018, 519.
- [32] J. Radlow, A two-dimensional singular integral equation of diffraction theory, *Bulletin of the American Mathematical Society*, (70)4 1964, 596-599.
- [33] Y. Saad, M.H. Schultz, GMRES: A generalized minimal residual algorithm for solving nonsymmetric linear system, *SIAM Journal on scientific and statistical computing*, 7(3) 1986, 856-869.
- [34] A. Soulaïmani, N.B. Salah, Y. Saad, Enhanced GMRES acceleration techniques for some CFD problems, *International Journal of Computational Fluid Dynamics*, 16(1) 2002, 1-20.
- [35] J. Shen, T. Tang, L. Wang, *Spectral Methods: Algorithms, Analysis and Applications*, Springer Publishing Company, Incorporated, 2011.
- [36] S.C. Shiralashetti, R.A. Mundewadi, S.S. Naregal, B. Veeresh, Haar Wavelet Collocation Method for the Numerical Solution of Nonlinear Volterra-Fredholm-Hammerstein Integral Equations, *Global Journal of Pure and Applied Mathematics*, 13(2) 2017, 463-474.
- [37] M.S. Tong, A Stable Integral Equation Solver for Electromagnetic Scattering by Large Scatterers with Concave Surface, *Progress In Electromagnetics Research*, 74 2007, 113-130.
- [38] E. Voltchkova, Integro-Differential Equations for Option Prices in Exponential Levy Models, *Finance and Stochastics*, 9(3) 2005, 299-325.
- [39] P. Wolfe, Eigenfunctions of the Integral Equation for the Potential of the Charged Disk, *Journal of Mathematical Physics*, 12(7) 1971, 1215-1218.
- [40] S. Yousefi, M. Razzaghi, Legendre wavelets method for the nonlinear VolterraFredholm integral equations, *Mathematics and computers in simulation*, 70(1) 2005, 1-8.