

# Mittag-Leffler-Hyer-Ulam-Rassias stability of cubic functional equation

V. Kalvandi,<sup>1</sup> M. E. Samei<sup>2 3</sup>

**Abstract:** In this paper, we prove the Mittag-Leffler-Hyers-Ulam-Rassias stability for cubic functional equation by using the fixed point alternative theorem. As a consequence, we show that the cubic multipliers are superstable under some conditions.

**Keywords:** Quadratic functional; Superstable; Cubic multipliers.

**2020 Mathematics Subject Classification:** 45M10, 65P40, 65L20, 70H14.

**Receive:** 26 June 2021, **Accepted:** 6 August 2021

## 1 Introduction

The stability theory for functional equations started with a problem related to the stability of group homomorphism that was considered by Ulam in 1940(see [27, 28]). Ulam considered the following question: “Let  $\mathcal{G}_1$  be a group and let  $\mathcal{G}_2$  be a group endowed with a metric  $\rho$ . Given  $\varepsilon > 0$ , does there exist a  $\delta > 0$  such that if a mapping  $\vartheta : \mathcal{G}_1 \rightarrow \mathcal{G}_2$  satisfies the inequality

$$\rho(\vartheta(y \cdot z), \vartheta(y)\vartheta(z)) < \delta,$$

for each  $y, z \in \mathcal{G}_1$ , then we can find a homomorphism  $\hat{h} : \mathcal{G}_1 \rightarrow \mathcal{G}_2$  such that

$$\rho(\vartheta(y), \hat{h}(y)) < \varepsilon,$$

for almost all  $y \in \mathcal{G}_1$ ?”. In other words, we are looking for situations when the homomorphisms are stable, i.e., if a mapping is almost a homomorphism, then there exists a true homomorphism near it. An affirmative answer to this question was given by Hyers in [12], for the case of Banach spaces. This answer, in this case, says that the Cauchy functional equation is stable in the sense of Hyers - Ulam. In 1950, T. Aoki was the second author to treat this problem for additive mapping (for more details, see [5, 9]). In 1978, Rassias generalized the theorem of Hyers by considering the stability problem with unbounded Cauchy differences ([20]). He has introduced a new type of stability which is called the Hyers - Ulam - Rassias stability. Research in stability theory is now very extensive and many papers and books have been published (for instance, consider [13, 17, 21, 23–25]). Recently, the Ulam stability of first order, second order and third order differential equations were investigated in a series of papers [3, 18, 29]. Murali and Selvan investigated the Mittag - Leffler - Hyers - Ulam stability and Mittag - Leffler - Hyers - Ulam - Rassias stability of a homogeneous and non-homogeneous linear differential equation of first order  $u'(t) + \ell u(t) = 0$  and  $u'(t) + \ell u(t) = r(t)$ , for all  $t \in I := [a, b]$ ,  $u(t), r(t) \in C(I)$ , by using the Laplace Transforms Methods [18].

The functional equation

$$u(y+z) + u(y-z) = 2u(y) + 2u(z), \quad (1.1)$$

is called quadratic functional equation. Also, every solution (for example  $u(y) = ay^2$ ) of functional Equation (1.1) is said to be a quadratic mapping. A Hyers - Ulam stability problem for the quadratic functional equation was proved by Skof for mappings  $u : \mathcal{N} \rightarrow \mathcal{B}$ , where  $\mathcal{N}$  is a normed space and  $\mathcal{B}$  is a Banach space ([26]). Jun *et al.* in [15] introduced the functional equation

$$u(2y+z) + u(2y-z) = 2u(y+z) + 2u(y-z) + 12u(y), \quad (1.2)$$

which is somewhat different from Eq. (1.1). It is easy to see that function  $u(y) = ay^3$  is a solution of Eq. (1.2). Thus, it is natural that Equation (1.2) is called a cubic functional equation and every solution of this cubic functional equation

<sup>1</sup>Department of Mathematics, Razi University, Kermanshah, Iran *E-mail addresses:* [Vida.kalvandi@yahoo.com](mailto:Vida.kalvandi@yahoo.com)

<sup>2</sup>Corresponding author: Department of Mathematics, Faculty of Basic Science, Bu-Ali Sina University, Hamedan, Iran, [mesamei@basu.ac.ir](mailto:mesamei@basu.ac.ir), [mesamei@gmail.com](mailto:mesamei@gmail.com)

<sup>3</sup>Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan

is said to be a cubic function. One year after that in [16], they solved the generalized Hyers - Ulam - Rassias stability of a cubic functional equation

$$u(y + 2z) + u(y - 2z) + 6u(y) = u(y + z) + 4u(y - z). \quad (1.3)$$

Recently, Bodaghi *et al.* in [6, 7, 19] proved the superstability of quadratic double centralizers and of quadratic multipliers on Banach algebras by fixed point methods. They studied  $n$ -variable mappings that are quartic in each variable, showed that the conditions defining such mappings can be unified in a single functional equation and by using an alternative fixed point method to prove the Hyers - Ulam stability for the multi-quartic functional equations in the normed spaces. In fact a mapping  $c : \mathcal{G}^n \rightarrow \mathcal{S}$  is called multicubic if it is cubic, i.e., it satisfies the cubic functional equation

$$c(2y + z) + c(2yz) = 2c(y + z) + 2c(yz) + 12c(y), \quad (1.4)$$

in each variable. where  $\mathcal{G}$  is a commutative group and  $\mathcal{S}$  is a linear space ([8]). Also the stability and super stability of cubic double centralizers of Banach algebras which are strongly without order had been established in [11] by Eshaghi Gordji *et al.*

In this paper, we investigate the Mittag - Leffler - Hyers - Ulam - Rassias stability (MLHUR-stability) by using the alternative fixed point for the cubic functional Equation (1.2) and their correspondent cubic multipliers.

## 2 Main results

Throughout this section,  $\mathcal{N}$  is a normed vector space and  $\mathcal{B}$  is a Banach space.

Now, for the given mapping  $u : \mathcal{N} \rightarrow \mathcal{B}$ , we consider the functional equation

$$\mathbb{D}[u](y, z) := u(2y + z) + u(2y - z) - 2u(y + z) - 2u(y - z) - 12u(y). \quad (2.1)$$

We need the following known fixed point theorem, which is useful for our results.

**Theorem 2.1.** [10] *(The fixed point alternative)* Suppose that  $(\Upsilon, \rho)$  be a complete generalized metric space. and let  $\mathfrak{J} : \Upsilon \rightarrow \Upsilon$  be a strictly contractive mapping with the Lipschitz constant  $\ell < 1$ . Then for each element  $y \in \Upsilon$ , either

$$\rho(\mathfrak{J}^n(y), \mathfrak{J}^{n+1}(y)) = \infty,$$

for all  $n \geq 0$ , or there exists a natural number  $N_0$  such that:

- (i) For all  $n \geq N_0$ ,  $\rho(\mathfrak{J}^n(y), \mathfrak{J}^{n+1}(y)) = \infty$ .
- (ii) The sequence  $\{\mathfrak{J}^n(y)\}$  is convergent to a fixed point  $y^*$  of  $\mathfrak{J}$ .
- (iii)  $y^*$  is the unique fixed point of  $\mathfrak{J}$  in

$$\Omega = \left\{ \omega \in \Upsilon : \rho(\mathfrak{J}^{N_0}(y), \omega) < \infty \right\}.$$

(iv) For all  $\omega \in \Omega$ ,

$$\rho(\omega, y^*) \leq \frac{1}{1 - \ell} \rho(\mathfrak{J}(\omega), \omega).$$

### 2.1 Stability of cubic function equations

**Definition 2.2.** The Mittag - Leffler function of one parameter is denoted by  $\mathcal{E}_\zeta(z)$  and defined as

$$\mathcal{E}_\zeta(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(k\zeta + 1)},$$

where  $z, \zeta \in \mathbb{C}$  with  $\text{Re}(\zeta) > 0$ .

**Theorem 2.3.** Let  $u : \mathcal{N} \rightarrow \mathcal{B}$  be a mapping with  $u(0) = 0$  and let  $\psi : \mathcal{N} \times \mathcal{N} \rightarrow [0, \infty)$  be a function satisfying

$$\lim_{n \rightarrow \infty} \frac{\psi(2^n y, 2^n z)}{8^n} = 0,$$

and

$$\|\mathbb{D}[u](y, z)\| \leq \psi(y, z)\mathcal{E}_\zeta(y), \quad (2.2)$$

for all  $y, z \in \mathcal{N}$ . If there exists  $\ell \in (0, 1)$  such that

$$\psi(2y, 0)\mathcal{E}_\zeta(2y) \leq 8\ell\psi(y, 0)\mathcal{E}_\zeta(y), \quad (2.3)$$

for each  $y \in \mathcal{N}$ , then there exists a unique cubic mapping  $\mathfrak{c} : \mathcal{N} \rightarrow \mathcal{B}$  such that for  $y \in \mathcal{N}$ ,

$$\|\mathfrak{u}(y) - \mathfrak{c}(y)\| \leq \frac{\ell}{2(1-\ell)}\psi(y, 0)\mathcal{E}_\zeta(y). \quad (2.4)$$

*Proof.* We consider the set

$$\Upsilon := \left\{ \mu : \mathcal{N} \rightarrow \mathcal{B} \mid \mu(0) = 0 \right\},$$

and introduce the generalized metric on  $\Upsilon$  as follows:

$$\rho(\mu_1, \mu_2) := \inf \left\{ C \in (0, \infty] : \|\mu_1(y) - \mu_2(y)\| \leq C\psi(y, 0)\mathcal{E}_\zeta(y), \forall y \in \mathcal{N} \right\},$$

where  $\mathcal{E}_\zeta$  is Mittag - Leffler function. If there is not exists such constant  $C$  then  $\rho(\mu_1, \mu_2) = \infty$ , otherwise. One can prove that the metric space  $(\Upsilon, \rho)$  is complete. Now, we define the mapping  $\mathcal{J} : \Upsilon \rightarrow \Upsilon$  by  $\mathcal{J}[\mu](y) = \frac{1}{8}\mu(2y)$ , for  $y \in \mathcal{X}$ . If  $\mu_1, \mu_2 \in \Upsilon$  such that  $\rho(\mu_1, \mu_2) < C$ , by definition of  $\rho$  and  $\mathcal{J}$ , we have

$$\left\| \frac{1}{8}\mu_1(2y) - \frac{1}{8}\mu_2(2y) \right\| \leq \frac{1}{8}C\psi(2y, 0)\mathcal{E}_\zeta(2y),$$

for each  $y \in \mathcal{N}$ . By using Eq. (2.3), we get

$$\left\| \frac{1}{8}\mu_1(2y) - \frac{1}{8}\mu_2(2y) \right\| \leq C\ell\psi(y, 0)\mathcal{E}_\zeta(y),$$

for all  $y \in \mathcal{N}$ . The above inequality shows that

$$\rho(\mathcal{J}[\mu_1], \mathcal{J}[\mu_2]) \leq \ell\rho(\mu_1, \mu_2),$$

for almost all  $\mu_1, \mu_2 \in \Upsilon$ . Hence,  $\mathcal{J}$  is a strictly contractive mapping on  $\Upsilon$  with Lipschitz constant  $\ell$ . Putting  $z = 0$  in inequality (2.2), using (2.3), and dividing both sides of the resulting inequality by 16, we have

$$\left\| \frac{1}{8}\mathfrak{u}(2y) - \mathfrak{u}(y) \right\| \leq \frac{1}{16}\psi(y, 0)\mathcal{E}_\zeta(y) \leq \frac{\ell}{2}\psi\left(\frac{y}{2}, 0\right)\mathcal{E}_\zeta\left(\frac{y}{2}\right),$$

for all  $y \in \mathcal{N}$ . Thus,  $\rho(\mathfrak{u}, \mathcal{J}[\mathfrak{u}]) \leq \frac{1}{2}\ell < \infty$ . By Theorem 2.1, the sequence  $\{\mathcal{J}[\mathfrak{u}]\}$  converges to a fixed point  $\mathfrak{c} : \mathcal{N} \rightarrow \mathcal{B}$  in the set

$$\Upsilon_1 = \{ \mu \in \Upsilon : \rho(\mathfrak{u}, \mu) < \infty \},$$

that is

$$\mathfrak{c}(y) = \lim_{n \rightarrow \infty} \frac{\mathfrak{u}(2^n y)}{8^n}. \quad (2.5)$$

By Theorem 2.1, we have

$$\rho(\mathfrak{u}, \mathfrak{c}) \leq \frac{\rho(\mathfrak{u}, \mathcal{J}[\mathfrak{u}])}{1-\ell} \leq \frac{\ell}{2(1-\ell)}. \quad (2.6)$$

It follows from inequality (2.6), that inequality (2.4) holds for all  $y \in \mathcal{X}$ . Substituting  $y, z$  by  $2^n y, 2^n z$  in (2.2), we have

$$\begin{aligned} \|\mathbb{D}[\mathfrak{c}](y, z)\| &= \lim_{n \rightarrow \infty} \frac{1}{8^n} \|\mathbb{D}[\mathfrak{u}](2^n y, 2^n z)\| \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{8^n} \psi(2^n y, 2^n z) \mathcal{E}_\zeta(2^n y) = 0, \end{aligned}$$

for all  $y \in \mathcal{N}$ . Therefore  $\mathfrak{c}$  is a cubic mapping which is unique.  $\square$

**Corollary 2.4.** Let  $\eta$  and  $\lambda$  be non-negative real numbers such that  $\eta < 3$ . Suppose that  $\mathfrak{u} : \mathcal{N} \rightarrow \mathcal{B}$  is a mapping satisfying

$$\|\mathbb{D}[\mathfrak{u}](y, z)\| \leq \lambda(\|y\|^\eta + \|z\|^\eta)\mathcal{E}_\zeta(y),$$

for all  $y, z \in \mathcal{N}$ . Then, there exists a unique cubic mapping  $\mathfrak{c} : \mathcal{N} \rightarrow \mathcal{B}$  such that

$$\|\mathfrak{u}(y) - \mathfrak{c}(y)\| \leq \frac{2^\eta \lambda}{2(8 - 2^\eta)} \|y\|^\eta \mathcal{E}_\zeta(y),$$

for all  $x \in \mathcal{N}$ .

*Proof.* The result follows from Theorem 2.3 by using  $\psi(y, z) = \lambda(\|y\|^\eta + \|z\|^\eta)$ .  $\square$

**Corollary 2.5.** *Let  $\eta_1, \eta_2, \lambda$  be non-negative real numbers such that  $\eta_1$  and  $\eta_2 \in (3, \infty)$ . Suppose a mapping  $u : \mathcal{N} \rightarrow \mathcal{B}$  satisfies*

$$\|\mathbb{D}[u](y, z)\| \leq \lambda\|y\|^{\eta_1}\|z\|^{\eta_2}\mathcal{E}_\zeta(y), \quad (2.7)$$

for all  $y, z \in \mathcal{N}$ . Then,  $u$  is cubic mapping on  $\mathcal{N}$ .

*Proof.* Letting  $y = z = 0$  in (2.7), we get  $u(0) = 0$ . Once more, if we put  $z = 0$  in (2.7), we have  $u(2y) = 8u(y)$  for all  $y \in \mathcal{N}$ . It is easy to see that by induction, we have  $u(2^n y) = 8^n u(y)$ , and so

$$u(y) = \frac{1}{8^n}u(2^n y),$$

for all  $y \in \mathcal{N}$  and  $n \in \mathbb{N}$ . Now, it follows from Theorem 2.3 that  $u$  is a cubic mapping.  $\square$

Note that in Corollary 2.5, if  $\eta_1 + \eta_2 \in (0, 3)$  and  $\eta_1 > 0$  such that the inequality (2.7) holds, then by applying

$$\psi(y, z) = \lambda\|y\|^{\eta_1}\|z\|^{\eta_2},$$

in Theorem 2.3,  $u$  is again a cubic be mapping.

**Theorem 2.6.** *Let  $u : \mathcal{N} \rightarrow \mathcal{B}$  be a mapping with  $u(0) = 0$ , and let  $\psi : \mathcal{N} \times \mathcal{N} \rightarrow [0, \infty)$  be a function satisfying*

$$\lim_{n \rightarrow \infty} 8^n \left( \frac{y}{2^n}, \frac{z}{2^n} \right) = 0, \quad (2.8)$$

and

$$\|\mathbb{D}[u](y, z)\| \leq \psi(y, z)\mathcal{E}_\zeta(y), \quad (2.9)$$

for each  $y, z \in \mathcal{N}$ . If there exists  $\ell \in (0, 1)$  such that

$$\psi(y, 0)\mathcal{E}_\zeta(y) \leq \frac{1}{8}\ell\psi(2y, 0)\mathcal{E}_\zeta(2y), \quad (2.10)$$

for almost all  $y \in \mathcal{N}$ , then there exists a unique cubic mapping  $c : \mathcal{N} \rightarrow \mathcal{B}$  such that

$$\|u(y) - c(y)\| \leq \frac{\ell}{16(1-\ell)}\psi(y, 0)\mathcal{E}_\zeta(y). \quad (2.11)$$

*Proof.* We consider the set

$$\Upsilon := \left\{ \mu : \mathcal{N} \rightarrow \mathcal{B} \mid \mu(0) = 0 \right\}$$

and introduce the generalized metric on  $\Upsilon$

$$\rho(\mu_1, \mu_2) := \inf \left\{ C \in (0, \infty) : \|\mu_1(y) - \mu_2(y)\| \leq C\psi(y, 0)\mathcal{E}_\zeta(y), \forall y \in \mathcal{N} \right\},$$

if there exists such constant  $C$  and  $\rho(\mu_1, \mu_2) = \infty$ , otherwise. It is easy to show that  $(\Upsilon, \rho)$  is complete. We will show that the mapping  $\mathcal{J} : \Upsilon \rightarrow \Upsilon$  define by

$$\mathcal{J}[\mu](y) = 8\mu\left(\frac{y}{2}\right),$$

for  $y \in \mathcal{N}$ , is strictly contractive. For given  $\mu_1, \mu_2 \in \Upsilon$  such that  $\rho(\mu_1, \mu_2) < C$ , we have

$$\left\| 8\mu_1\left(\frac{y}{2}\right) - 8\mu_2\left(\frac{y}{2}\right) \right\| \leq C\psi\left(\frac{y}{2}, 0\right)\mathcal{E}_\zeta\left(\frac{y}{2}\right),$$

for all  $y \in \mathcal{N}$ . By using inequality (2.10), we obtain

$$\left\| 8\mu_1\left(\frac{y}{2}\right) - 8\mu_2\left(\frac{y}{2}\right) \right\| \leq C\ell\psi(y, 0)\mathcal{E}_\zeta(y),$$

for all  $y \in \mathcal{N}$ . It follows from the last inequality that  $\rho(\mathcal{J}[\mu_1], \mathcal{J}[\mu_2]) \leq \ell\rho(\mu_1, \mu_2)$  for all  $\mu_1, \mu_2 \in \Upsilon$ . Hence,  $\mathcal{J}$  is a strictly contractive mapping on  $\Upsilon$  with Lipschitz constant  $\ell$ . By putting  $z = 0$  and replacing  $y$  by  $\frac{y}{2}$  in inequality (2.9) and using inequality (2.10), then by dividing both sides of the resulting inequality by 2, we have

$$\left\| 8u\left(\frac{y}{2}\right) - u(y) \right\| \leq \frac{1}{2}\psi\left(\frac{y}{2}, 0\right)\mathcal{E}_\zeta\left(\frac{y}{2}\right) \leq \frac{1}{16}\ell\psi(y, 0)\mathcal{E}_\zeta(y),$$

for all  $y \in \mathcal{N}$ . Hence,

$$\rho(u, \mathcal{J}[u]) \leq \frac{L}{16} < \infty.$$

By applying the fixed point alternative, there exists a unique mapping  $\mathfrak{v} : \mathcal{N} \rightarrow \mathcal{B}$  in the set  $\Upsilon_1 = \{\mu \in \Upsilon : \rho(u, \mu) < \infty\}$ , such that

$$\mathfrak{c}(y) = \lim_{n \rightarrow \infty} 8^n u\left(\frac{y}{2^n}\right), \quad (2.12)$$

for every  $y \in \mathcal{N}$ . Again, Theorem 2.1 shows that

$$\rho(u, \mathfrak{c}) \leq \frac{\rho(u, \mathcal{J}[u])}{1 - \ell} \leq \frac{\ell}{16(1 - \ell)}, \quad (2.13)$$

where the inequality (2.13) implies the relation (2.11). Replacing  $y, z$  by  $2^n y, 2^n z$  in (2.9), respectively, and using (2.8) and (2.12), we conclude

$$\|\mathbb{D}[\mathfrak{c}](y, z)\| = \lim_{n \rightarrow \infty} 8^n \left\| \mathbb{D}[u]\left(\frac{y}{2^n}, \frac{z}{2^n}\right) \right\| \leq \lim_{n \rightarrow \infty} 8^n \psi\left(\frac{y}{2^n}, \frac{z}{2^n}\right) = 0,$$

for all  $y, z \in \mathcal{N}$ . Therefore  $\mathfrak{c}$  is a cubic mapping.  $\square$

## 2.2 Stability of cubic multipliers

In this section, we investigate the Mittag - Leffler - Hyers - Ulam stability and the superstability of cubic multipliers.

**Definition 2.7.** A cubic multiplier on an algebra  $\mathcal{G}$  is a cubic mapping  $\mathfrak{T} : \mathcal{G} \rightarrow \mathcal{G}$  such that

$$y\mathfrak{T}(z) = \mathfrak{T}(y)z,$$

for each  $y, z \in \mathcal{G}$ .

The following theorem introduces a cubic multipliers on Banach algebras.

**Theorem 2.8.** Let  $u : \mathcal{G} \rightarrow \mathcal{G}$  be a mapping with  $u(0) = 0$  and let  $\psi : \mathcal{G}^4 \rightarrow [0, \infty)$  be a function such that

$$\|\mathbb{D}[u](y, z) + u(v)w - vu(w)\| \leq \psi(y, z, v, w)\mathcal{E}_\zeta(y), \quad (2.14)$$

for all  $y, z, v, w \in \mathcal{G}$ . If there exists a constant  $\ell \in (0, 1)$  such that

$$\psi(2y, 2z, 2v, 2w) \leq 8\ell\psi(y, z, v, w), \quad (2.15)$$

for all  $y, z, v, w \in \mathcal{G}$ , then there exists a unique cubic multiplier  $\mathfrak{T}$  on  $\mathcal{G}$  satisfying

$$\|u(y) - \mathfrak{T}(y)\| \leq \frac{\ell}{2(1 - \ell)}\psi(y, y, 0, 0)\mathcal{E}_\zeta(y), \quad (2.16)$$

for each  $y \in \mathcal{X}$ .

*Proof.* It follows from the relation (2.15) that

$$\lim_{n \rightarrow \infty} \frac{\psi(2^n y, 2^n z, 2^n v, 2^n w)}{8^n} = 0, \quad (2.17)$$

for all  $y, z, v, w \in \mathcal{G}$ . Putting  $z = v = w = 0$  in (2.14), we obtain

$$\|2u(2y) - 16u(y)\| \leq \psi(y, 0, 0, 0)\mathcal{E}_\zeta(y),$$

for every  $y \in \mathcal{G}$ . Thus,

$$\left\| \frac{1}{8}u(2y) - u(y) \right\| \leq \frac{1}{16}\psi(y, 0, 0, 0)\mathcal{E}_\zeta(y), \quad (2.18)$$

for all  $y \in \mathcal{G}$ . Now similar to the proof of theorem in previous section, we consider the set

$$\Upsilon := \{\mu : \mathcal{G} \rightarrow \mathcal{G} \mid \mu(0) = 0\},$$

and introduce the generalized metric on  $\Upsilon$  as:

$$\rho(\mu_1, \mu_2) := \inf \{C \in \mathbb{R}^+ : \|\mu_1(y) - \mu_2(y)\| \leq C\psi(y, 0, 0, 0)\mathcal{E}_\zeta(y), \forall y \in \mathcal{G}\},$$

if there exists such constant  $C$ , and  $\rho(\mu_1, \mu_2) = \infty$ , otherwise. The metric space  $(\Upsilon, \rho)$  is complete, and by the same reasoning as in the proof of Theorem 2.3, the mapping  $\Phi : \Upsilon \rightarrow \Upsilon$  defined by

$$\Phi[\mu](y) = \frac{1}{8}\mu(2y),$$

for  $y \in \mathcal{G}$  is strictly contractive on  $\mathcal{X}$  and has a unique fixed point  $\mathfrak{T}$  such that

$$\lim_{n \rightarrow \infty} \rho(\Phi^n \mathbf{u}, \mathfrak{T}) = 0,$$

i.e.,

$$\mathfrak{T}(y) = \lim_{n \rightarrow \infty} \frac{\mathbf{u}(2^n y)}{8^n}, \quad (2.19)$$

for all  $y \in \mathcal{G}$ . By Theorem 2.3 shows that  $\mathfrak{T}$  is a cubic mapping. If we substitute  $v$  and  $w$  by  $2^n v$  and  $2^n w$  in (2.14), respectively, and put  $y = z = 0$  and we divide the both sides of the obtained inequality by  $2^{4n}$ , we get

$$\left\| v \frac{1}{8^n} \mathbf{u}(2^n w) - \frac{1}{8^n} \mathbf{u}(2^n v) w \right\| = 0.$$

Passing to the limit as  $n \rightarrow \infty$  and from (2.17), we conclude that  $v\mathfrak{T}(w) = \mathfrak{T}(v)w$  for all  $v, w \in \mathcal{G}$ .  $\square$

**Corollary 2.9.** *Let  $\eta, \lambda$  be non-negative real numbers with  $\eta < 3$  and let  $\mathbf{u} : \mathcal{G} \rightarrow \mathcal{G}$  be a mapping with  $\mathbf{u}(0) = 0$  such that*

$$\|\mathbb{D}[\mathbf{u}](y, z) + \mathbf{u}(v)w - v\mathbf{u}(w)\| \leq \lambda(\|y\|^\eta + \|z\|^\eta + \|v\|^\eta + \|w\|^\eta)\mathcal{E}_\zeta(y),$$

for all  $y, z, v, w \in \mathcal{G}$ . Then, there exists a unique cubic multiplier  $\mathfrak{T}$  on  $\mathcal{G}$  satisfying

$$\|\mathbf{u}(y) - \mathfrak{T}(y)\| \leq \frac{2^{\eta-1}\lambda}{8-2^\eta}\|y\|^\eta,$$

for all  $y \in \mathcal{G}$ .

*Proof.* The proof follows from Theorem 2.8 by taking

$$\psi(y, z, v, w) = \lambda(\|y\|^\eta + \|z\|^\eta + \|v\|^\eta + \|w\|^\eta),$$

for all  $y, z, v, w \in \mathcal{G}$ .  $\square$

**Corollary 2.10.** *Let  $\eta_j$  for  $j = 1, 2, 3, 4$ ,  $\lambda$  be non-negative real numbers with*

$$\sum_{j=1}^4 \eta_j < 3,$$

and let  $\mathbf{u}$  maps  $\mathcal{G}$  to  $\mathcal{G}$  be a mapping with  $\mathbf{u}(0) = 0$  such that

$$\|\mathbb{D}[\mathbf{u}](y, z) + \mathbf{u}(v)w - v\mathbf{u}(w)\| \leq \lambda(\|y\|^{\eta_1} + \|z\|^{\eta_2} + \|v\|^{\eta_3} + \|w\|^{\eta_4})\mathcal{E}_\zeta(y),$$

for all  $y, z, v, w \in \mathcal{G}$ . Then,  $\mathbf{u}$  is a cubic multiplier on  $\mathcal{G}$ .

*Proof.* It is enough to let

$$\psi(y, z, v, w) = \lambda(\|y\|^{\eta_1} + \|z\|^{\eta_2} + \|v\|^{\eta_3} + \|w\|^{\eta_4}),$$

in Theorem 2.8.  $\square$

### 3 Conclusion

Quadratic functional equation was used to characterize inner product spaces [1, 4, 14]. Several other functional equations were also to characterize inner product spaces. A square norm on an inner product space satisfies the important parallelogram equality

$$\|y + z\|^2 + \|y - z\|^2 = 2(\|y\|^2 + \|z\|^2).$$

This research has made an attempt to analyse the MLHU stability and the MLHUR stability of linear differential equation with constant coefficients. Also we have showed that the Mittag - Leffler function plays an immodest role to prove the stability of differential equation. This new method of stability unifies different classes of differential equations, which may inspire further research in this domain

## Declarations

### Availability of Data and Materials

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

### Competing interests

The authors declare that they have no competing interests.

### Funding

Not applicable.

### Authors' contributions

The authors declare that the study was realized in collaboration with equal responsibility. All authors read and approved the final manuscript.

## References

- [1] J. Aczél, J. Dhombres, *Functional Equations in Several Variables*, Cambridge University Press, 1989.
- [2] M. Akkouchi, Hyers-Ulam-Rassias Stability of nonlinear Volterra integral equations via a fixed point approach, *Acta. Univ. Apulen. Math. Inform.*, 26 2011, 257-266.
- [3] Q.H. Alqifiary, J. K. Miljanovic, Note on the stability of system of differential equations  $\dot{x}(t) = f(t, x(t))$ , *Gen*, 20(1) 2014, 27-33.
- [4] D. Amir, *Characterizations of Inner Product Spaces*, Birkhuser, Basel, 1986.
- [5] T. Aoki, On the stability of the linear transformation in Banach spaces, *Journal of the mathematical society of Japan*, 2(1-2) (1950), 64-66.
- [6] A. Bodaghi, I.A. Alias, M. Eshaghi Gordji, On the stability of quadratic double centralizers and quadratic multipliers: a fixed point approach, *Journal of Inequalities and Applications*, 2011(1) 2011, 1-9.
- [7] A. Bodaghi, I.A. Alias, M.H. Ghahramani, Approximately cubic functional equations and cubic multipliers, *Journal of Inequalities and Applications*, 2011(1) 2011, 1-8.
- [8] A. Bodaghi, C. Park, O.T Mewomo, Multiquartic functional equations, *Advances in Difference Equations*, 2019(1) 2019, 1-10.
- [9] D.G. Bourgin, Approximately isometric and multiplicative transformations on continuous function rings, *Duke Mathematical Journal*, 16(2) 1949, 385-397.
- [10] J.B. Diaz, B. Margolis, A fixed point theorem of the alternative for contractions on a generalized complete metric space, *Bulletin of the American Mathematical Society*, 74(2) 1968, 305-309
- [11] M. Eshdaghi Gordji, S. Shams, M. Ramezani, A. Ebadian, Approximately cubic double centralizers, *Nonlinear Func Anal Appl*, 15(3) 2010, 503-512.
- [12] D.H. Hyers, On the stability of the linear functional equation, *Proceedings of the national academy of sciences of the United States of America*, 27(4) 1941, 222-224.
- [13] D.H. Hyers, G. Isac, T.M. Rassias, *Stability of Functional Equation in Several Variables*, Rirkhäuser, Basel, 1998.
- [14] P. Jordan, J. Von Neumann, On inner products in linear metric spaces, *Annals of Mathematics*, 36(3) 1935, 719-723.
- [15] K.W. Jun, H.M. Kim, The generalized Hyers - Ulam - Rassias stability of a cubic functional equation, *Journal of Mathematical Analysis and Applications*, 274(2) 2002, 867-878.
- [16] K.W. Jun, H.M. Kim, On the Hyers - Ulam - Rassias stability of a general cubic functional equation, *Mathematical Inequalities and Applications*, 6(2) 2003, 289-302.
- [17] V. Kalvandi, M.E. Samei, New stability results for a sum-type fractional  $q$ -integro-differential equation, *J. Adv. Math. Stud*, 12(2) 2019, 201-209.
- [18] R. Murali, A. Ponmana Selvan, Mittag-LefflerHyersUlam stability of a linear differential equations of first order using Laplace transforms, *Canad. J. Appl. Math.*, 2(2) 2020, 47-59.
- [19] C. Park, A. Bodaghi, Two multi-cubic functional equations and some results on the stability in modular spaces, *Journal of Inequalities and Applications* 2020(1) 2020, 1-16.

- [20] T.M. Rassias, On the stability of the linear mapping in Banach spaces, *Proceedings of the American mathematical society* 72(2) 1978, 297-300
- [21] T.M. Rassias, The problem of S.M. Ulam for approximately multiplicative mappings, *Journal of Mathematical Analysis and Applications*, 246(2) 2000, 352-378.
- [22] T.M. Rassias, On the stability of functional equations in Banach spaces, *Journal of Mathematical Analysis and Applications* 251(1) 2000, 264-284.
- [23] T.M. Rassias, On the stability of functional equations and a problem of Ulam, *Acta Applicandae Mathematica*, 62(1) 2000, 23-130.
- [24] T.M. Rassias, *Functional Equations, Inequalities and Applications*, Kluwer Academic, Dordrecht, Boston and London, 2003.
- [25] M.E. Samei, V. Hedayati, Sh. Rezapour, Existence results for a fraction hybrid differential inclusion with caputo-hadamard type fractional derivative, *Advances in Difference Equations*, 2019(1) 2019, 1-15.
- [26] F. Skof, Propriet locall e approssimazione di operatori. *Rend sem Math Fis Milano, Rendiconti del Seminario Matematico e Fisico di Milano*, 53(1) 1983, 113-129.
- [27] S.M. Ulam, *Problem in Modern Mathematics*, Chapter VI, Science Editions, Wiley, New York, 1960.
- [28] S.M. Ulam, *A Collection of the Mathematical Problems*, Interscience Publ. New York, 1960.
- [29] J. Xue, Hyers-Ulam stability of linear differential equations of second order with constant coefficient, *talian Journal of Pure and Applied Mathematics*, 32 2014, 419-424.